

## SYSTEMATIZING PROBABILITY:

$$S = \{s_1, s_2, \dots, s_n\} \leftarrow \text{The } s_i \text{ are outcomes.}$$

$$\begin{array}{c} \swarrow \quad \downarrow \quad \searrow \\ w(s_1) \quad w(s_2) \quad w(s_n) \end{array} \leftarrow \text{The } w(s_i) \text{ are numbers } \geq 0.$$

ERASE



1



$$S = \{s_1, s_2, \dots, s_n\} \leftarrow \text{The } s_i \text{ are outcomes.}$$

$$\begin{array}{c} \swarrow \quad \downarrow \quad \searrow \\ w(s_1) \quad w(s_2) \quad w(s_n) \end{array} \leftarrow \text{The } w(s_i) \text{ are numbers } \geq 0.$$

If  $E = \{s_1, s_3, s_7\} \subset S$ , then  $\Pr[E] = w(s_1) + w(s_3) + w(s_7)$ .  $\Pr[E]$  is the probability that  $E$  will occur. Note that

$$1) 0 \leq \Pr[E] \leq 1$$

since  $w(s_1) + w(s_2) + \dots + w(s_n) = 1$ . Also

$$2) \Pr[S] = 1 \text{ AND } \Pr[\emptyset] = 0.$$

If  $E$  and  $F$  are disjoint ( $E \cap F = \emptyset$ ), then

$$3) \Pr[E \cup F] = \Pr[E] + \Pr[F]$$

This last statement follows from common sense ...

ERASE



2



Lecture 9

$$E = \{s_1, s_5, s_7, s_{22}\}$$

$$F = \{s_4, s_{59}, s_{78}, s_{212}, s_{21244}\}$$

$$w(s_1) + w(s_5) + w(s_7) + w(s_{22})$$

$$w(s_4) + w(s_{59}) + w(s_{78}) + w(s_{212}) + w(s_{21244})$$

$\Pr[E]$

$\Pr[F]$

$$\{s_1, s_5, s_7, s_{22}, s_4, s_{59}, s_{78}, s_{212}, s_{21244}\}$$

$$w(s_1) + w(s_5) + w(s_7) + w(s_{22}) + w(s_4) + w(s_{59}) + w(s_{78}) + w(s_{212}) + w(s_{21244})$$

$\Pr[E] + \Pr[F]$

ERASE



3



ERASE



4



$$\{s_1, s_5, s_7, s_{22}, s_4, s_{59}, s_{78}, s_{212}, s_{21244}\}$$

$$w(s_1) + w(s_5) + w(s_7) + w(s_{22}) + w(s_4) + w(s_{59}) + w(s_{78}) + w(s_{212}) + w(s_{21244})$$

$\Pr[E] + \Pr[F]$

Clearly, this is also  $\Pr[E \cup F]$

$$\{s_1, s_5, s_7, s_{22}, s_4, s_{59}, s_{78}, s_{212}, s_{21244}\}$$

$$w(s_1) + w(s_5) + w(s_7) + w(s_{22}) + w(s_4) + w(s_{59}) + w(s_{78}) + w(s_{212}) + w(s_{21244})$$

$\Pr[E] + \Pr[F]$

Clearly, this is also  $\Pr[E \cup F]$

So,  $\Pr[E \cup F] = \Pr[E] + \Pr[F]$  ( $E \cap F = \emptyset$ )

ERASE



5



ERASE



6



**COMMENT 1:** The book adopts 1), 2), and 3) as axioms and then goes on to show that everything you need to know can be derived from these axioms. We will also derive what we need from 1), 2), and 3).

**COMMENT 2:** One immediate consequence of 3) follows by considering  $E \subset S$ :

$$E \cup E^c = S \quad E \cap E^c = \emptyset.$$

So by 3),

$$1 = \Pr[S] = \Pr[E \cup E^c] = \Pr[E] + \Pr[E^c]$$

or

$$1 = \Pr[E] + \Pr[E^c] \text{ for all } E \subset S$$

ERASE



7



**RECALL:** If  $A$  and  $B$  are sets, then

$$n(A \cup B) = n(A) + n(B) - n(A \cap B)$$

The only fact that we used in deriving this was:

$$n(A \cup B) = n(A) + n(B) \text{ for disjoint sets.}$$

We now have the corresponding statement holding true for probabilities

$$\Pr[A \cup B] = \Pr[A] + \Pr[B] \text{ for disjoint sets, so it}$$

should be true that

$$\Pr[A \cup B] = \Pr[A] + \Pr[B] - \Pr[A \cap B] \text{ for any sets.}$$


The reasoning is the same: Just label the segments of the Venn diagram with their probabilities -

ERASE




8





ERASE

9






So now use  $\Pr[E \cup F] = \Pr[E] + \Pr[F]$  for disjoint sets:

ERASE

10


Lecture 9



Noting that the two sets,  and , shown above are disjoint.


ERASE

11



ERASE

12



$$\Pr[A \cup B] = \Pr[A] - \Pr[A \cap B] + \Pr[A \cap B] + \Pr[B] - \Pr[A \cap B]$$

$$= \Pr[A] + \Pr[B] - \Pr[A \cap B]$$

ERASE

13

- All together:
- 1)  $0 \leq \Pr[E] \leq 1$
  - 2)  $\Pr[S] = 1$  AND  $\Pr[\emptyset] = 0$ .
  - 3)  $\Pr[E \cup F] = \Pr[E] + \Pr[F] - \Pr[E \cap F]$

ERASE

14

Minor comment:

Note that  $\Pr[\emptyset] = 0$  actually follows from  $\Pr[S] = 1$  and the fact  $1 = \Pr[E] + \Pr[E^c]$  (applied to  $\emptyset$  and its complement  $S$ ).

**EXAMPLE:** Suppose  $E$  and  $F$  are events in a sample space  $S$  with  $\Pr[E] = .65$ ,  $\Pr[F] = .4$  and  $\Pr[E \cup F] = .75$ . Find

- a)  $\Pr[E \cap F]$
- b)  $\Pr[G]$  where  $G$  is the set of all outcomes which are in exactly one of the sets  $E$  and  $F$ .

ERASE

15

**EXAMPLE:** Suppose  $E$  and  $F$  are events in a sample space  $S$  with  $\Pr[E] = .65$ ,  $\Pr[F] = .4$  and  $\Pr[E \cup F] = .75$ . Find

- a)  $\Pr[E \cap F]$
- b)  $\Pr[G]$  where  $G$  is the set of all outcomes which are in exactly one of the sets  $E$  and  $F$ .

a)  $\Pr[E \cup F] = \Pr[E] + \Pr[F] - \Pr[E \cap F]$

ERASE

16

**EXAMPLE:** Suppose  $E$  and  $F$  are events in a sample space  $S$  with  $\Pr[E] = .65$ ,  $\Pr[F] = .4$  and  $\Pr[E \cup F] = .75$ . Find

- $\Pr[E \cap F]$
- $\Pr[G]$  where  $G$  is the set of all outcomes which are in exactly one of the sets  $E$  and  $F$ .

a)  $\Pr[E \cup F] = \Pr[E] + \Pr[F] - \Pr[E \cap F]$   
 $.75 = .65 + .4 - \Pr[E \cap F]$

ERASE



17



**EXAMPLE:** Suppose  $E$  and  $F$  are events in a sample space  $S$  with  $\Pr[E] = .65$ ,  $\Pr[F] = .4$  and  $\Pr[E \cup F] = .75$ . Find

- $\Pr[E \cap F]$
- $\Pr[G]$  where  $G$  is the set of all outcomes which are in exactly one of the sets  $E$  and  $F$ .

a)  $\Pr[E \cup F] = \Pr[E] + \Pr[F] - \Pr[E \cap F]$   
 $.75 = .65 + .4 - \Pr[E \cap F]$   
 $.75 = 1.05 - \Pr[E \cap F]$   
 $\Pr[E \cap F] = .3$

ERASE



18



Lecture 9

**EXAMPLE:** Suppose  $E$  and  $F$  are events in a sample space  $S$  with  $\Pr[E] = .65$ ,  $\Pr[F] = .4$  and  $\Pr[E \cup F] = .75$ . Find

- $\Pr[E \cap F] = .3$
- $\Pr[G]$  where  $G$  is the set of all outcomes which are in exactly one of the sets  $E$  and  $F$ .

ERASE




19



**EXAMPLE:** Suppose  $E$  and  $F$  are events in a sample space  $S$  with  $\Pr[E] = .65$ ,  $\Pr[F] = .4$  and  $\Pr[E \cup F] = .75$ . Find

- $\Pr[E \cap F] = .3$
- $\Pr[G]$  where  $G$  is the set of all outcomes which are in exactly one of the sets  $E$  and  $F$ .

Venn Diagram:



ERASE



20



**EXAMPLE:** Suppose  $E$  and  $F$  are events in a sample space  $S$  with  $\Pr[E] = .65$ ,  $\Pr[F] = .4$  and  $\Pr[E \cup F] = .75$ . Find

- $\Pr[E \cap F] = .3$
- $\Pr[G]$  where  $G$  is the set of all outcomes which are in exactly one of the sets  $E$  and  $F$ .


ERASE



21




Venn Diagram:



**EXAMPLE:** Suppose  $E$  and  $F$  are events in a sample space  $S$  with  $\Pr[E] = .65$ ,  $\Pr[F] = .4$  and  $\Pr[E \cup F] = .75$ . Find

- $\Pr[E \cap F] = .3$
- $\Pr[G]$  where  $G$  is the set of all outcomes which are in exactly one of the sets  $E$  and  $F$ .

Venn Diagram:



ERASE



22



**EXAMPLE:** Suppose  $E$  and  $F$  are events in a sample space  $S$  with  $\Pr[E] = .65$ ,  $\Pr[F] = .4$  and  $\Pr[E \cup F] = .75$ . Find

- $\Pr[E \cap F] = .3$
- $\Pr[G]$  where  $G$  is the set of all outcomes which are in exactly one of the sets  $E$  and  $F$ .


ERASE



23




Venn Diagram:



**EXAMPLE:** Suppose  $E$  and  $F$  are events in a sample space  $S$  with  $\Pr[E] = .65$ ,  $\Pr[F] = .4$  and  $\Pr[E \cup F] = .75$ . Find

- $\Pr[E \cap F] = .3$
- $\Pr[G]$  where  $G$  is the set of all outcomes which are in exactly one of the sets  $E$  and  $F$ .

Venn Diagram:



ERASE




24



**EXAMPLE:** Suppose E and F are events in a sample space S with  $\Pr[E] = .65$ ,  $\Pr[F] = .4$  and  $\Pr[E \cup F] = .75$ . Find

- $\Pr[E \cap F] = .3$
- $\Pr[G]$  where G is the set of all outcomes which are in exactly one of the sets E and F.

Venn Diagram:



ERASE



25




Lecture 9

**EXAMPLE:** Suppose E and F are events in a sample space S with  $\Pr[E] = .65$ ,  $\Pr[F] = .4$  and  $\Pr[E \cup F] = .75$ . Find

- $\Pr[E \cap F] = .3$
- $\Pr[G]$  where G is the set of all outcomes which are in exactly one of the sets E and F.

Venn Diagram:



ERASE



26



**EXAMPLE:** A widget manufacturer make widgets. The widgets are often mismanufactured (too large or too small) and/or mislabeled (incorrect label or correct label in wrong place). The probabilities for a randomly selected widget selected from the production line are shown below:

ERASE



27



**EXAMPLE:** A widget manufacturer make widgets. The widgets are often mismanufactured (too large or too small) and/or mislabeled (incorrect label or correct label in wrong place). The probabilities for a randomly selected widget selected from the production line are shown below:

ERASE



28



Added note: Not clearly stated in problem, but we will use it,  $H_1$  and  $H_2$  are disjoint.

	EVENT	Probability
F	Acceptable	.72
G	$G_1$ Too large	.12
	$G_2$ Too small	.08
H	$H_1$ Incorrect label	.15
	$H_2$ Correct label/wrong place	.02

	EVENT	Probability
F	Acceptable	.72
G	$G_1$ Too large	.12
	$G_2$ Too small	.08
H	$H_1$ Incorrect label	.15
	$H_2$ Correct label/wrong place	.02

**EXAMPLE:** For randomly selected widget, find:

- Probability widget is mismanufactured
- Probability widget is not mismanufactured
- Probability widget is mislabeled

ERASE



29



	EVENT	Probability
F	Acceptable	.72
$G_1$	Too large	.12
	Too small	.08
$H_1$	Incorrect label	.15
	Correct label/wrong place	.02

$G = G_1 \cup G_2$

$H = H_1 \cup H_2$

**EXAMPLE:** For randomly selected widget, find:

- Probability widget is mismanufactured  $\Pr[G]$
- Probability widget is not mismanufactured  $\Pr[G^c]$
- Probability widget is mislabeled  $\Pr[H]$

ERASE



30



	EVENT	Probability
F	Acceptable	.72
$G_1$	Too large	.12
	Too small	.08
$H_1$	Incorrect label	.15
	Correct label/wrong place	.02

$G = G_1 \cup G_2$

$H = H_1 \cup H_2$

**EXAMPLE:** For randomly selected widget, find:

- Probability widget is mismanufactured  $\Pr[G]$
- Probability widget is not mismanufactured  $\Pr[G^c]$
- Probability widget is mislabeled  $\Pr[H]$

ERASE



31



$\Pr[\text{mismanufactured}] = \Pr[G] = \Pr[G_1 \cup G_2]$

	EVENT	Probability
F	Acceptable	.72
$G_1$	Too large	.12
	Too small	.08
$H_1$	Incorrect label	.15
	Correct label/wrong place	.02

$G = G_1 \cup G_2$   
disjoint

$H = H_1 \cup H_2$

**EXAMPLE:** For randomly selected widget, find:

- Probability widget is mismanufactured  $\Pr[G]$
- Probability widget is not mismanufactured  $\Pr[G^c]$
- Probability widget is mislabeled  $\Pr[H]$

ERASE



32



$\Pr[\text{mismanufactured}] = \Pr[G] = \Pr[G_1 \cup G_2] = \Pr[G_1] + \Pr[G_2]$   
 $= .12 + .08 = .20$

$G_1 \cap G_2 = \emptyset$

	EVENT	Probability
F	Acceptable	.72
$G_1$	Too large	.12
	Too small	.08
$H_1$	Incorrect label	.15
	Correct label/wrong place	.02

$G = G_1 \cup G_2$   
disjoint

$H = H_1 \cup H_2$

**EXAMPLE:** For randomly selected widget, find:

- Probability widget is mismanufactured  $\Pr[G] = .20$
- Probability widget is not mismanufactured  $\Pr[G^c]$
- Probability widget is mislabeled  $\Pr[H]$

EVENT		Probability
F	Acceptable	.72
$G_1$	Too large	.12
$G_2$	Too small	.08
$H_1$	Incorrect label	.15
$H_2$	Correct label/wrong place	.02

$G = G_1 \cup G_2$   
 $H = H_1 \cup H_2$

ERASE



33



Lecture 9

**EXAMPLE:** For randomly selected widget, find:

- Probability widget is mismanufactured  $\Pr[G] = .20$
- Probability widget is not mismanufactured  $\Pr[G^c]$
- Probability widget is mislabeled  $\Pr[H]$

$\Pr[\text{NOT mismanufactured}] = \Pr[G^c] = 1 - \Pr[G] = 1 - .20 = .80$

EVENT		Probability
F	Acceptable	.72
$G_1$	Too large	.12
$G_2$	Too small	.08
$H_1$	Incorrect label	.15
$H_2$	Correct label/wrong place	.02

$G = G_1 \cup G_2$   
 $H = H_1 \cup H_2$

ERASE



34



**EXAMPLE:** For randomly selected widget, find:

- Probability widget is mismanufactured  $\Pr[G] = .20$
- Probability widget is not mismanufactured  $\Pr[G^c] = .80$
- Probability widget is mislabeled  $\Pr[H]$

EVENT		Probability
F	Acceptable	.72
$G_1$	Too large	.12
$G_2$	Too small	.08
$H_1$	Incorrect label	.15
$H_2$	Correct label/wrong place	.02

$G = G_1 \cup G_2$   
 $H = H_1 \cup H_2$

ERASE



35



**EXAMPLE:** For randomly selected widget, find:

- Probability widget is mismanufactured  $\Pr[G] = .20$
- Probability widget is not mismanufactured  $\Pr[G^c] = .80$
- Probability widget is mislabeled  $\Pr[H]$

$\Pr[\text{mislabeled}] = \Pr[H] = \Pr[H_1 \cup H_2] = \Pr[H_1] + \Pr[H_2]$   
 $= .15 + .02 = .17$

$H_1 \cap H_2 = \emptyset$

EVENT		Probability
F	Acceptable	.72
$G_1$	Too large	.12
$G_2$	Too small	.08
$H_1$	Incorrect label	.15
$H_2$	Correct label/wrong place	.02

$G = G_1 \cup G_2$   
 $H = H_1 \cup H_2$   
*disjoint*

ERASE



36



**EXAMPLE:** For randomly selected widget, find:

- Probability widget is mismanufactured  $\Pr[G] = .20$
- Probability widget is not mismanufactured  $\Pr[G^c] = .80$
- Probability widget is mislabeled  $\Pr[H] = .17$

EVENT		Probability
F	Acceptable	.72
$G_1$	Too large	.12
$G_2$	Too small	.08
$H_1$	Incorrect label	.15
$H_2$	Correct label/wrong place	.02

$G = G_1 \cup G_2$   
 $H = H_1 \cup H_2$

ERASE



37



**EXAMPLE:** For randomly selected widget, find:

- Probability widget is mismanufactured  $\Pr[G] = .20$
- Probability widget is not mismanufactured  $\Pr[G^c] = .80$
- Probability widget is mislabeled  $\Pr[H] = .17$

EVENT		Probability
F	Acceptable	.72

ERASE



38



**EXAMPLE:** For randomly selected widget, find:

- Probability widget is mismanufactured  $\Pr[G] = .20$
- Probability widget is not mismanufactured  $\Pr[G^c] = .80$
- Probability widget is mislabeled  $\Pr[H] = .17$
- Probability widget is either mislabeled or mismanufactured
- Probability widget is both mislabeled and mismanufactured

EVENT		Probability
F	Acceptable	.72

ERASE



39



**EXAMPLE:** For randomly selected widget, find:

- Probability widget is mismanufactured  $\Pr[G] = .20$
- Probability widget is not mismanufactured  $\Pr[G^c] = .80$
- Probability widget is mislabeled  $\Pr[H] = .17$
- Probability widget is either mislabeled or mismanufactured  $\Pr[G \cup H] = \Pr[F^c]$
- Probability widget is both mislabeled and mismanufactured  $\Pr[G \cap H]$

EVENT		Probability
F	Acceptable	.72

ERASE



40



**EXAMPLE:** For randomly selected widget, find:

- a) Probability widget is mismanufactured  $\Pr[G] = .20$
- b) Probability widget is not mismanufactured  $\Pr[G^c] = .80$
- c) Probability widget is mislabeled  $\Pr[H] = .17$
- d) Probability widget is either mislabeled or mismanufactured  $\Pr[G \cup H] = \Pr[F^c]$
- e) Probability widget is both mislabeled and mismanufactured  $\Pr[G \cap H]$

$$\Pr[F^c] = 1 - \Pr[F] = 1 - .72 = .28$$

EVENT	Probability
F Acceptable	.72

ERASE



41



**EXAMPLE:** For randomly selected widget, find:

- a) Probability widget is mismanufactured  $\Pr[G] = .20$
- b) Probability widget is not mismanufactured  $\Pr[G^c] = .80$
- c) Probability widget is mislabeled  $\Pr[H] = .17$
- d) Probability widget is either mislabeled or mismanufactured  $\Pr[G \cup H] = \Pr[F^c] = .28$
- e) Probability widget is both mislabeled and mismanufactured  $\Pr[G \cap H]$

EVENT	Probability
F Acceptable	.72

ERASE



42



Lecture 9

**EXAMPLE:** For randomly selected widget, find:

- a) Probability widget is mismanufactured  $\Pr[G] = .20$
- b) Probability widget is not mismanufactured  $\Pr[G^c] = .80$
- c) Probability widget is mislabeled  $\Pr[H] = .17$
- d) Probability widget is either mislabeled or mismanufactured  $\Pr[G \cup H] = \Pr[F^c] = .28$
- e) Probability widget is both mislabeled and mismanufactured  $\Pr[G \cap H]$

$$\Pr[G \cup H] = \Pr[G] + \Pr[H] - \Pr[G \cap H] \Rightarrow$$

EVENT	Probability
F Acceptable	.72

ERASE



43



**EXAMPLE:** For randomly selected widget, find:

- a) Probability widget is mismanufactured  $\Pr[G] = .20$
- b) Probability widget is not mismanufactured  $\Pr[G^c] = .80$
- c) Probability widget is mislabeled  $\Pr[H] = .17$
- d) Probability widget is either mislabeled or mismanufactured  $\Pr[G \cup H] = \Pr[F^c] = .28$
- e) Probability widget is both mislabeled and mismanufactured  $\Pr[G \cap H]$

$$\Pr[G \cup H] = \Pr[G] + \Pr[H] - \Pr[G \cap H] \Rightarrow$$

$$.28 = .20 + .17 - \Pr[G \cap H] \Rightarrow \Pr[G \cap H] = .09$$

EVENT	Probability
F Acceptable	.72

ERASE



44



ERASE



45



ERASE



46



ERASE



47



ERASE



48

